

Reference: PA0025

Technical Data Sheet

Product: Sabouraud Dextrose Agar

Specification

Solid medium for the enumeration and cultivation of fungi according to the Pharmacopeial Harmonised Method and ISO standard.

re				

Storage **Shelf Life** 20 Prepared Plates **Packaging Details** 90 mm 1 box with 2 packs of 10 plates/pack. Single 3 months 2-14 °C cellophane.. with: 21 ± 2 ml

Composition

Composition (g/l):	
D(+)-Glucose	40.0
Peptone from casein	5.00
Meat Peptone	
Agar	
3.	

Description / Technique

Sabouraud Dextrose Agar is a modification of the classical Sabouraud medium for the cultivation of fungi. This new formula helps to maintain the morphology of fungi, providing a reliable medium for both cultivation and differentiation.

Its selectivity is due to a low pH and a high glucose concentration, which together with incubation at a relatively lower temperature (20 -25°C) favours the growth of fungi while discouraging that of bacteria.

The mixture of peptones employed has been selected to provide the fungi with all their nitrogen requirements.

Spread the plate streaking methodology or by spiral method.

Each laboratory must evaluate the results according to their specifications.

Quality control

Physical/Chemical control

Color: Straw-coloured yellow pH: 5.6 ± 0.2 at 25°C

Microbiological control

Growth Promotion Test according to harmonized pharmacopoeial monographs and test methods & ISO 11133:2014/A1:2018

Spiral Spreading: Practical range 50 - 100 CFU (productivity).

Aerobiosis. Incubation at 20-25°C. Reading ≤5 days.

Microorganism	Growth
Candida albicans ATCC® 10231, WDCM 00054	Good (≥70 %)
Aspergillus brasiliensis ATCC® 16404, WDCM 00053	Good (≥70 %)
S.cerevisiae ATCC® 9763, WDCM 00058	Good (≥70 %)
Sterility control	

Incubation 48 hours at 30-35 °C and 48 hours at 20-25 °C: NO GROWTH.

Check at 7 days after incubation in same conditions.

Page 1 / 2 Revision date: 08/04/21



Reference: PA0025

Technical Data Sheet

Product: Sabouraud Dextrose Agar

Bibliography

·AJELLO, L. (1957) Cultural Methods for Human Pathogenic Fungi J. Chron. Dis. 5:545-551.

·COLIPA (1997) Guidelines on Microbial Quality Management (MQM). Brussels.

·EUROPEAN PHARMACOPOEIA 10.0 (2020) 10th ed. § 2.6.13. Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. EDQM. Council of Europe. Strasbourg.

GEORGE, L.K., AJELLO, L. & PAPAGEORGE, C. (1954) Use of Cycloheximide in the Selective Isolation of Fungi Pathogenic to Man. J. Lab. Clin. Med, 44 (422-428).

·HANTSCHKE, D. (1968) Mykosen, 11, (769-778).

. ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.

·ISO 16212 Standard (2017) Cosmetics - Microbiology - Enumeration of yeast and mould.

PAGANO, J. LEVIN, J.D. and TREJO, W. (1957-58) Diagnostic Medium for Differentiation of Species of Candida. Antibiotics Annual, 137

·SABOURAUD, R. (1910) Les Teignes. Masson, Paris.

·USP 33 - NF 28 (2011) <62> Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. USP Corp. Inc. Rockville. MD. USA.

Page 2 / 2 Revision date: 08/04/21